

## Quiz for Wave Medical Clinical Rotation Companion

**1) What age group has the peak incidence of Alopecia Areata?**

- a) < 5 years
- b) 5-14 years
- c) 15-29 years
- d) 30-50 years
- e) > 50 years

**2) What does the abbreviation LAD stand for?**

- a) Left Anterior Descending Artery
- b) Left Axis Deviation
- c) Lymphadenopathy
- d) All of the above
- e) None of the above

**3) How many types of Salter Harris fractures are listed in the Clinical Anatomy section?**

- a) 6
- b) 7
- c) 5
- d) 2
- e) 4

**4) Premature Atrial Contraction is a type of which of the following?**

- a) Congestive Heart Failure
- b) Heart Blockage
- c) Premature Labor
- d) Irregular Heart Beat
- e) None of the Above

---

**5) Which of the following drug combinations should not be used if at all avoidable?**

- a) Kanamycin and Selenium
- b) Medrol and Penicillin
- c) Baridium and Teicoplanin
- d) Teicoplanin and Kanamycin
- e) None of the above

**6) Torsades is a \_\_\_\_\_ type of Rhythm Strip?**

- a) Atrial
- b) Ventricular
- c) Heart blocks
- d) Irregular beats
- e) Pacemaker

**7) Under Clinical Rotations, how many subtopics are in the Surgery section ?**

- a) 5
- b) 9
- c) 16
- d) 14
- e) 10

**8) Which of the following combinations results in the most severe interaction?**

- a) Hydrocodone and Pirbuterol
- b) Morphine and Paxil
- c) Hydrocodone and Morphine
- d) Morphine and Pirbuterol
- e) No interactions found between any of the listed combinations

---

**9) The Extra Cellular Fluid equation is a part of which class of Medical Calculator?**

- a) Body Mass Index
- b) Body Weight, Ideal
- c) Body Water, Total
- d) Body Weight, Adjusted
- e) All of the above

**10) What is not an indication of Ventilator Management (Basic Principles section)?**

- a) Ventilatory Failure
- b) Oxygenation Failure
- c) Mechanical Failure
- d) Vital Capacity
- e) Positive End Expiratory Pressure

**11) Which of the following is not an adverse drug reaction to Accutane?**

- a) Cheilitis
- b) Thinning hair
- c) Hypertriglyceridemia
- d) All of the above
- e) None of the above

**12) How many different dosing options are in the renal impairment section in administering Milrinone?**

- a) 5
- b) 2
- c) 7
- d) 6
- e) 3

---

13) What condition does Auspitz's Sign suggest?

- a) Blowfish Poisoning
- b) HELLP Syndrome
- c) Psoriasis
- d) Reiter's Syndrome
- e) Onychomycosis

14) Which of the following is not a caution when administering Milrinone?

- a) Atrial fib/flutter
- b) Electrolyte abnormalities
- c) Short-term use only
- d) Antidepressants
- e) Renal impairment

15) Which type of Salter-Harris fracture crosses through the epiphyseal plate?

- a) Type I, "S"
- b) Type II, "A"
- c) Type III, "L"
- d) Type IV, "T"
- e) Type V, "ER"

16) What is the definitive test for Herpes Simplex Virus Type 2 infection?

- a) Clinical exam
- b) Serology
- c) Culture and typing
- d) Tzanck smear
- e) CSF testing

17) What is the first step in administering Milrinone?

- a) 50 mcg/kg IV over 20 minutes
- b) 60 mcg/kg IV over 20 minutes
- c) 65 mcg/kg IV over 10 minutes
- d) 60 mcg/kg IV over 10 minutes
- e) 50 mcg/kg IV over 10 minutes

## Answers

- 1) c
- 2) d
- 3) c
- 4) d
- 5) d
- 6) b
- 7) d
- 8) b
- 9) c
- 10) e
- 11) d
- 12) d
- 13) c
- 14) d
- 15) d
- 16) a
- 17) e