

Quiz for Wave Medical Primary Pro

- 1) Out of the drugs listed, which combination should never be used because of its serious, potential life threatening interaction?
- a) Glimepiride and benazepril
 - b) Glimepiride and suprefact
 - c) Glimepiride and ethanol
 - d) Glimepiride and clotrimazole
 - e) None of the Above
- 2) In the Evidence-Based Medicine Table of Contents, how many entries are there in the neurology section that are in regards to Bell's Palsy?
- a) 1
 - b) 2
 - c) 3
 - d) 4
 - e) None of the Above
- 3) What is the mean corpuscular volume of if a patient has an 80% hemocrit level over 10 red blood cells?
- a) 70 fL
 - b) 80 fL
 - c) 90 fL
 - d) 100 fL
 - e) None of the Above
- 4) Out of the drugs listed, which combination should never be used because of its serious, potential life threatening interaction?
- a) Azithromycin and bemiparin
 - b) Azithromycin and dicumarol
 - c) Azithromycin and protamine
 - d) Azithromycin and pimoziide
 - e) None of the Above

5) Which is not a differential diagnosis that is possible for contact dermatitis?

- a) Atopic dermatitis
- b) Psoriasis
- c) Acne rosacea
- d) Dermatophytes
- e) None of the Above

6) Which is not an adverse reaction to diphenhydramine?

- a) Decreased cognitive function in geriatrics
- b) Fainting
- c) Sedation
- d) Confusion
- e) None of the Above

7) According to the glossary, "Raccoon Eyes" are seen in?

- a) LASIK complications
- b) Basilar skull fractures
- c) Conjunctivitis
- d) Glaucoma
- e) None of the Above

8) The abbreviation JVD stands for?

- a) Juvenile Vascular Dissension
- b) Jugular Venous Access
- c) Jugular Venous Distension
- d) Junctional Vascular Tachycardia
- e) None of the Above

9) The drug Meloxicam is classified into what pregnancy category?

- a) A
- b) B
- c) C
- d) D
- e) E

10) What percent of patients may experience dizziness as an adverse reaction to lamotrigine?

- a) 25%
- b) 35%
- c) 38%
- d) 48%
- e) None of the Above

11) A patient measures 164.4 mEq/L of sodium and 100.7 mg/dL of glucose, what is there corrected sodium?

- a) 164.41 mEq/L
- b) 123.34 mEq/L
- c) 16.41 mEq/L
- d) 20.23 mEq/L
- e) None of the Above

12) Out of the drugs listed, which combination should never be used because of its serious, potential life threatening interaction?

- a) Ganciclovir and ampicillin
- b) Ganciclovir and zidovudine
- c) Ganciclovir and nafcillin
- d) Ganciclovir and cloxacillin
- e) None of the Above

13) What is the recommended paediatric dosing for glatiramer?

- a) 20 mg SC qD
- b) 40 mg SC qD
- c) 2 mg SC qD
- d) Not recommended
- e) None of the Above

14) The acronym for remembering transient causes of urinary incontinence in the elderly is?

- a) DOPRS
- b) DAPRRS
- c) DAPPER
- d) DIAPPRS
- e) None of the Above

15) The recommended adult dose of montelukast is?

- a) 1 mg PO BID
- b) 10 mg PO qHS
- c) 4 mg PRN
- d) 1.5 mg PO qHS
- e) None of the Above

Answers

- 1) c
- 2) a
- 3) b
- 4) d
- 5) c
- 6) b
- 7) b
- 8) c
- 9) c
- 10) c
- 11) a
- 12) b
- 13) d
- 14) d
- 15) b